

Stable functorial equivalence of blocks

Serge Bouc and Deniz Yılmaz

Abstract

Let k be an algebraically closed field of characteristic $p > 0$, let R be a commutative ring and let \mathbb{F} be an algebraically closed field of characteristic 0. We introduce the category $\overline{\mathcal{F}}_{Rpp_k}^\Delta$ of stable diagonal p -permutation functors over R . We prove that the category $\overline{\mathcal{F}}_{\mathbb{F}pp_k}^\Delta$ is semisimple and give a parametrization of its simple objects in terms of the simple diagonal p -permutation functors.

We also introduce the notion of a stable functorial equivalence over R between blocks of finite groups. We prove that if G is a finite group and if b is a block idempotent of kG with an abelian defect group D and Frobenius inertial quotient E , then there exists a stable functorial equivalence over \mathbb{F} between the pairs (G, b) and $(D \rtimes E, 1)$.

Keywords: block, diagonal p -permutation functors, functorial equivalence, Frobenius inertial quotient.

MSC2020: 16S34, 20C20, 20J15.

1 Introduction

In past decades, various notions of equivalences between blocks of finite groups have been studied such as splendid Morita equivalence, splendid Rickard equivalence, p -permutation equivalence, isotypies and perfect isometries ([Br90], [BX08], [BP20]). These equivalences are related to prominent conjectures in modular representation theory such as Broué's abelian defect group conjecture (Conjecture 9.7.6 in [L18]), Puig's finiteness conjecture (Conjecture 6.4.2 in [L18]) and Donovan's conjecture (Conjecture 6.1.9 in [L18]).

Recently, in [BY22] we introduced another equivalence of blocks, namely *functorial equivalence*, using the notion of diagonal p -permutation functors: Let k be an algebraically closed field of characteristic $p > 0$, let \mathbb{F} be an algebraically closed field of characteristic 0 and let R be a commutative ring. We denote by Rpp_k^Δ the category whose objects are finite groups and for finite groups G and H whose morphisms from H to G are the Grothendieck group $RT^\Delta(G, H)$ of diagonal p -permutation (kG, kH) -bimodules. An R -linear functor from Rpp_k^Δ to ${}_R\mathbf{Mod}$ is called a *diagonal p -permutation functor*. To each pair (G, b) of a finite group G and a block idempotent

b of kG , we associate a canonical diagonal p -permutation functor over R , denoted by $RT_{G,b}^\Delta$. If (H, c) is another such pair, we say that (G, b) and (H, c) are *functorially equivalent over R* if the functors $RT_{G,b}^\Delta$ and $RT_{H,c}^\Delta$ are isomorphic.

In [BY22] we proved that the category of diagonal p -permutation functors over \mathbb{F} is semisimple, parametrized simple functors and provided three equivalent descriptions of the decomposition of the functor $\mathbb{F}T_{G,b}^\Delta$ in terms of the simple functors ([BY22, Corollary 6.15 and Theorem 8.22]). We proved that the number of isomorphism classes of simple modules, the number of ordinary irreducible characters, and the defect groups are preserved under functorial equivalences over \mathbb{F} ([BY22, Theorem 10.5]). Moreover we proved that for a given finite p -group D , there are only finitely many pairs (G, b) , where G is a finite group and b is a block idempotent of kG , up to functorial equivalence over \mathbb{F} ([BY22, Theorem 10.6]) and we provided a sufficient condition for two blocks to be functorially equivalent over \mathbb{F} in the situation of Broué's abelian defect group conjecture ([BY22, Theorem 11.1]).

In this paper, we introduce the notion of *stable* diagonal p -permutation functors and *stable* functorial equivalences. We denote by $\overline{Rpp_k^\Delta}$ the quotient category of Rpp_k^Δ by the morphisms that factor through the trivial group. A *stable diagonal p -permutation functor over R* is an R -linear functor from $\overline{Rpp_k^\Delta}$ to $R\text{Mod}$, or equivalently, a diagonal p -permutation functor which vanishes at the trivial group. In particular, the simple diagonal p -permutation functors $S_{L,u,V}$ with $L \neq 1$ are (simple) stable diagonal p -permutation functors. Our first main result is the following.

1.1 Theorem *The category $\overline{\mathcal{F}_{\mathbb{F}pp_k}^\Delta}$ of stable diagonal p -permutation functors over \mathbb{F} is semisimple. The simple stable diagonal p -permutation functors are precisely the simple diagonal p -permutation functors $S_{L,u,V}$ with $L \neq 1$.*

Given a finite group G and a block idempotent b of kG , we define a stable diagonal p -permutation functor $\overline{RT_{G,b}^\Delta}$ similar to $RT_{G,b}^\Delta$, see Definition 4.1. Note that $\overline{RT_{G,b}^\Delta}$ is the zero functor if and only if b has defect 0. We say that two pairs (G, b) and (H, c) are *stably functorially equivalent over R* if the functors $\overline{RT_{G,b}^\Delta}$ and $\overline{RT_{H,c}^\Delta}$ are isomorphic. For a block algebra kGb , let $k(kGb)$ and $l(kGb)$ denote the number of irreducible ordinary characters and the number of irreducible Brauer characters of b , respectively.

1.2 Theorem *Let b be a block idempotent of kG and let c be a block idempotent of kH .*

- (i) *The pairs (G, b) and (H, c) are stably functorially equivalent over \mathbb{F} if and only if the multiplicities of $S_{L,u,V}$ in $\mathbb{F}T_{G,b}^\Delta$ and $\mathbb{F}T_{H,c}^\Delta$ are the same for any simple diagonal p -permutation functor $S_{L,u,V}$ with $L \neq 1$. In this case, (G, b) and (H, c) are functorially equivalent over \mathbb{F} if and only if $l(kGb) = l(kHc)$.*
- (ii) *If the pairs (G, b) and (H, c) are stably functorially equivalent over \mathbb{F} , then b and c have isomorphic defect groups and one has*

$$k(kGb) - l(kGb) = k(kHc) - l(kHc).$$

We also consider the blocks with abelian defect groups and Frobenius inertial quotient.

1.3 Theorem *Let G be a finite group, b a block idempotent of kG with a nontrivial abelian defect group D . Let $E = N_G(D, e_D)/C_G(D)$ denote the inertial quotient of b . Suppose that E acts freely on $D \setminus \{1\}$. Then:*

- (i) *There exists a functorial equivalence over \mathbb{F} between (G, b) and $(D \rtimes E, 1)$ if and only if $l(kGb) = l(k(D \rtimes E))$.*
- (ii) *Suppose that E is abelian. Then there exists a functorial equivalence over \mathbb{F} between (G, b) and $(D \rtimes E, 1)$ if and only if (G, b) and $(D \rtimes E, 1)$ are p -permutation equivalent.*

In Section 2 we recall diagonal p -permutation functors and functorial equivalences between blocks. In Section 3 we introduce the category of stable diagonal p -permutation functors and prove Theorem 1.1. In Section 4 we introduce the notion of stable functorial equivalences between blocks and prove Theorem 1.2. Finally, in Section 5 we prove Theorem 1.3.

2 Preliminaries

(a) Let (P, s) be a pair where P is a p -group and s is a generator of a p' -group acting on P . We write $P\langle s \rangle := P \rtimes \langle s \rangle$ for the corresponding semi-direct product. We say that two pairs (P, s) and (Q, t) are *isomorphic* and write $(P, s) \cong (Q, t)$, if there is a group isomorphism $f : P\langle s \rangle \rightarrow Q\langle t \rangle$ that sends s to a conjugate of t . We set $\text{Aut}(P, s)$ to be the group of the automorphisms of the pair (P, s) and $\text{Out}(P, s) = \text{Aut}(P, s)/\text{Inn}(P\langle s \rangle)$. Recall from [BY20] that a pair (P, s) is called a D^Δ -pair, if $C_{\langle s \rangle}(P) = 1$.

(b) Let G and H be finite groups. We denote by $T(G)$ the Grothendieck group of p -permutation kG -modules and by $T^\Delta(G, H)$ the Grothendieck group of p -permutation (kG, kH) -bimodules whose indecomposable direct summands have twisted diagonal vertices. Let Rpp_k^Δ denote the following category:

- objects: finite groups.
- $\text{Mor}_{Rpp_k^\Delta}(G, H) = R \otimes_{\mathbb{Z}} T^\Delta(H, G) = RT^\Delta(H, G)$.
- composition is induced from the tensor product of bimodules.
- $\text{Id}_G = [kG]$.

An R -linear functor from Rpp_k^Δ to ${}_R\mathbf{Mod}$ is called a *diagonal p -permutation functor* over R . Together with natural transformations, diagonal p -permutation functors form an abelian category $\mathcal{F}_{Rpp_k}^\Delta$.

(c) Recall from [BY22] that the category $\mathcal{F}_{\mathbb{F}pp_k}^\Delta$ is semisimple. Moreover, the simple diagonal p -permutation functors, up to isomorphism, are parametrized by the isomorphism classes of

triples (L, u, V) where (L, u) is a D^Δ -pair, and V is a simple $\mathbb{F}\text{Out}(L, u)$ -module (see [BY22, Sections 6 and 7] for more details on simple functors).

(d) Let G be a finite group and b a block idempotent of kG . Recall from [BY22] that the block diagonal p -permutation functor $RT_{G,b}^\Delta$ is defined as

$$\begin{aligned} RT_{G,b}^\Delta : Rpp_k^\Delta &\rightarrow {}_R\text{Mod} \\ H &\mapsto RT^\Delta(H, G) \otimes_{kG} kGb. \end{aligned}$$

See [BY22, Section 8] for the decomposition of $\mathbb{F}T_{G,b}^\Delta$ in terms of the simple functors $S_{L,u,V}$.

(e) Let b be a block idempotent of kG and let c be a block idempotent of kH . We say that the pairs (G, b) and (H, c) are *functorially equivalent* over R , if the corresponding diagonal p -permutation functors $RT_{G,b}^\Delta$ and $RT_{H,c}^\Delta$ are isomorphic in $\mathcal{F}_{Rpp_k}^\Delta$ ([BY22, Definition 10.1]). By [BY22, Lemma 10.2] the pairs (G, b) and (H, c) are functorially equivalent over R if and only if there exists $\omega \in bRT^\Delta(G, H)c$ and $\sigma \in cRT^\Delta(H, G)b$ such that

$$\omega \cdot_H \sigma = [kGb] \quad \text{in} \quad bRT^\Delta(G, G)b \quad \text{and} \quad \sigma \cdot_G \omega = [kHc] \quad \text{in} \quad cRT^\Delta(H, H)c.$$

3 Stable diagonal p -permutation functors

In this section we introduce the category of stable diagonal p -permutation functors.

For a finite group G , let $P(G)$ denote the subgroup of $T(G)$ generated by the indecomposable projective kG -modules. Let also $\overline{T(G)}$ denote the quotient group $T(G)/P(G)$. For $X \in T(G)$, we denote by \overline{X} the image of X in $\overline{T(G)}$. If H is another finite group, we define $P(G, H)$ and $\overline{T^\Delta(G, H)}$ similarly.

3.1 Lemma *For finite groups G and H one has $P(G, H) = T^\Delta(G, 1) \circ T^\Delta(1, H)$*

Proof This follows from the fact that the projective indecomposable $k(G \times H)$ -modules are of the form $P \otimes_k Q$ where P and Q are projective indecomposable kG and kH -modules, respectively. \square

3.2 Definition Let $\overline{Rpp_k^\Delta}$ denote the following category:

- objects: finite groups.
- $\text{Mor}_{\overline{Rpp_k^\Delta}}(G, H) = R \otimes_{\mathbb{Z}} \overline{T^\Delta(H, G)} = \overline{RT^\Delta(H, G)}$.
- composition is induced from the tensor product of bimodules.
- $\text{Id}_G = \overline{[kG]}$.

3.3 Definition An R -linear functor $\overline{Rpp_k^\Delta} \rightarrow {}_R\mathbf{Mod}$ is called a *stable diagonal p -permutation functor* over R . Together with natural transformations, stable diagonal p -permutation functors form an abelian category $\overline{\mathcal{F}_{Rpp_k}^\Delta}$.

3.4 Remark The functor

$$\Gamma : \overline{\mathcal{F}_{Rpp_k}^\Delta} \rightarrow \mathcal{F}_{Rpp_k}^\Delta$$

obtained by composition with the projection $Rpp_k^\Delta \rightarrow \overline{Rpp_k^\Delta}$ gives a description of $\overline{\mathcal{F}_{Rpp_k}^\Delta}$ as a full subcategory of $\mathcal{F}_{Rpp_k}^\Delta$. Moreover, Γ has a left adjoint Σ , constructed as follows: If F is a diagonal p -permutation functor over R and G is a finite group, set

$$\overline{F}(G) := F(G)/RT^\Delta(G, \mathbf{1})F(\mathbf{1}).$$

Then \overline{F} is a diagonal p -permutation functor, equal to the quotient of F by the subfunctor generated by $F(\mathbf{1})$. Obviously, \overline{F} vanishes at the trivial group, so it is a stable diagonal p -permutation functor. The functor $\Sigma : F \mapsto \overline{F}$ is a left adjoint to the above functor Γ . In particular, $\overline{\mathcal{F}_{Rpp_k}^\Delta}$ is a reflective subcategory of $\mathcal{F}_{Rpp_k}^\Delta$.

Let G be a finite group. Recall that by [BY22, Corollary 8.23(i)], the multiplicity of the simple diagonal p -permutation functor $S_{1,1,\mathbb{F}}$ in the representable functor $\mathbb{F}T^\Delta(-, G)$ is equal to the number $l(kG)$ of the isomorphism classes of simple kG -modules. Let $\mathcal{I}(-, G)$ denote the sum of simple subfunctors of $\mathbb{F}T^\Delta(-, G)$ isomorphic to $S_{1,1,\mathbb{F}}$. Let also $\mathbb{F}\text{Proj}(-, G)$ denote the subfunctor of $\mathbb{F}T^\Delta(-, G)$ sending a finite group H to $\mathbb{F}\text{Proj}(H, G)$.

3.5 Lemma *The subfunctors $\mathcal{I}(-, G)$ and $\mathbb{F}\text{Proj}(-, G)$ of the representable functor $\mathbb{F}T^\Delta(-, G)$ are equal.*

Proof For finite groups G and H , the number of isomorphism classes of projective indecomposable $k(G \times H)$ -modules, or equivalently, the number of isomorphism classes of simple $k(G \times H)$ -modules is equal to the number of conjugacy classes of p' -elements of $G \times H$. Hence the \mathbb{F} -dimension of the evaluation $\mathbb{F}\text{Proj}(H, G)$ is equal to

$$l(k(G \times H)) = l(kG)l(kH)$$

which is equal to the \mathbb{F} -dimension of $l(kG)S_{1,1,\mathbb{F}}(H)$ by [BY22, Corollary 8.23(i)], and hence to the \mathbb{F} -dimension of $\mathcal{I}(H, G)$.

Note that $\mathbb{F}\text{Proj}(-, G)$ is equal to the functor

$$\mathbb{F}T^\Delta(-, 1) \circ \mathbb{F}T^\Delta(1, G).$$

Moreover $S_{L,u,V}(1) = 0$ for $L \neq 1$, and hence $\mathbb{F}T^\Delta(1, G) = \mathcal{I}(1, G)$. Therefore, one has

$$\mathbb{F}\text{Proj}(-, G) = \mathbb{F}T^\Delta(-, 1) \circ \mathbb{F}T^\Delta(1, G) = \mathbb{F}T^\Delta(-, 1) \circ \mathcal{I}(1, G) \subseteq \mathcal{I}(-, G).$$

Since the \mathbb{F} -dimensions of $\mathbb{F}\text{Proj}(H, G)$ and $\mathcal{I}(H, G)$ are the same for any finite group H , it follows that $\mathbb{F}\text{Proj}(-, G) = \mathcal{I}(-, G)$. \square

Proof of Theorem 1.1: For a finite group G , the representable diagonal p -permutation functor $\mathbb{F}T^\Delta(-, G)$ decomposes as a direct sum of simple functors $S_{L,u,V}$, and hence we have

$$\mathbb{F}T^\Delta(-, G) \cong \mathcal{I}(-, G) \bigoplus_{\substack{(L,u,V) \\ L \neq 1}} S_{L,u,V}^{m_{L,u,V}},$$

for some nonnegative integers $m_{L,u,V}$, where (L, u, V) runs over a set of isomorphism classes of D^Δ -pairs (L, u) with $L \neq 1$, and simple $\overline{\mathbb{F}\text{Out}(L, u)}$ -modules V . By Lemma 3.5, the representable stable diagonal p -permutation functor $\overline{\mathbb{F}T^\Delta}(-, G)$ is isomorphic to the direct sum

$$\bigoplus_{\substack{(L,u,V) \\ L \neq 1}} S_{L,u,V}^{m_{L,u,V}}$$

of simple diagonal p -permutation functors, and each of these simple functors is a simple stable diagonal p -permutation functor. Since the functor category $\overline{\mathcal{F}_{\mathbb{F}pp_k}^\Delta}$ is generated by the representable functors the result follows. \square

4 Stable functorial equivalences

Let G and H be finite groups.

4.1 Definition Let b a block idempotent of kG . The stable diagonal p -permutation functor $\overline{RT_{G,b}^\Delta}$ is defined as

$$\begin{aligned} \overline{RT_{G,b}^\Delta} : \overline{Rpp_k^\Delta} &\rightarrow {}_R\text{Mod} \\ H &\mapsto \overline{RT^\Delta(H, G) \otimes_{kG} kGb}. \end{aligned}$$

See Section 2(d) for the definition of $RT_{G,b}^\Delta$ and note that $\overline{RT_{G,b}^\Delta} = \Sigma \left(RT_{G,b}^\Delta \right)$.

4.2 Definition Let b be a block idempotent of kG and let c be a block idempotent of kH . We say that the pairs (G, b) and (H, c) are *stably functorially equivalent* over R , if their corresponding stable diagonal p -permutation functors $\overline{RT_{G,b}^\Delta}$ and $\overline{RT_{H,c}^\Delta}$ are isomorphic in $\overline{\mathcal{F}_{Rpp_k}^\Delta}$.

4.3 Lemma Let b be a block idempotent of kG and let c be a block idempotent of kH .

(a) The pairs (G, b) and (H, c) are stably functorially equivalent over R if and only if there exists $\omega \in bRT^\Delta(G, H)c$ and $\sigma \in cRT^\Delta(H, G)b$ such that

$$\omega \cdot_H \sigma = [kGb] + [P] \quad \text{in} \quad bRT^\Delta(G, G)b \quad \text{and} \quad \sigma \cdot_G \omega = [kHc] + [Q] \quad \text{in} \quad cRT^\Delta(H, H)c$$

for some $P \in R\text{Proj}(kGb, kGb)$ and $Q \in R\text{Proj}(kHc, kHc)$.

(b) If the pairs (G, b) and (H, c) are functorially equivalent over R , then they are also stably functorially equivalent over R .

Proof By the Yoneda lemma, the pairs (G, b) and (H, c) are stably functorially equivalent over R if and only if there exists $\bar{\omega} \in \overline{bRT^\Delta(G, H)c}$ and $\bar{\sigma} \in \overline{cRT^\Delta(H, G)b}$ such that

$$\bar{\omega} \cdot_H \bar{\sigma} = \overline{[kGb]} \quad \text{in} \quad \overline{bRT^\Delta(G, G)b} \quad \text{and} \quad \bar{\sigma} \cdot_G \bar{\omega} = \overline{[kHc]} \quad \text{in} \quad \overline{cRT^\Delta(H, H)c}.$$

Hence (a) follows and (b) is clear. \square

Proof of Theorem 1.2: (i) The first statement follows from Theorem 1.1 and the second statement follows since by [BY22, Corollary 8.23(i)] the multiplicity of the simple functor $S_{1,1,\mathbb{F}}$ in $\mathbb{F}T_{G,b}^\Delta$ is equal to $l(kGb)$.

(ii) The first statement follows from the proof of [BY22, Theorem 10.5(iii)] and the second statement follows from the proof of [BY22, Theorem 10.5(ii)]. \square

5 Blocks with Frobenius inertial quotient

(a) Recall the assumptions of Theorem 1.3: Let G be a finite group, b a block idempotent of kG with a nontrivial abelian defect group D . Let (D, e_D) be a maximal b -Brauer pair and let $E = N_G(D, e_D)/C_G(D)$ denote the inertial quotient of b . Suppose that E acts freely on $D \setminus \{1\}$. This condition is equivalent to requiring $D \rtimes E$ be a Frobenius group. Let \mathcal{F}_b be the fusion system of b with respect to (D, e_D) . Then \mathcal{F}_b is equal to the fusion system $\mathcal{F}_{D \rtimes E}(D)$ on D determined by $D \rtimes E$.

Under these assumptions, we know from [L18, Theorem 10.5.1] that there is a stable equivalence of Morita type between (G, b) and $(D \rtimes E, 1)$, induced by a bimodule with an endopermutation source. Since D is abelian, by [R96, Theorem 7.2] this bimodule admits an endosplit p -permutation resolution, which yields a stable p -permutation equivalence between (G, b) and $(D \rtimes E, 1)$, see for instance [L18, Remark 9.11.7]. In particular (G, b) and $(D \rtimes E, 1)$ are stably functorially equivalent.

Hereafter we give an alternative proof of the existence of this stable functorial equivalence, which relies only on Theorem 1.1. Together with Theorem 1.2, this now implies Part (i) of Theorem 1.3. Part (ii) of Theorem 1.3 follows from Part (i) and [L18, Theorem 10.5.10].

(b) Let $S_{L,u,V}$ be a simple diagonal p -permutation functor such that L is nontrivial and isomorphic to a subgroup of D . Recall that by [BY22, Theorem 8.22] the multiplicity of $S_{L,u,V}$ in $\mathbb{F}T_{G,b}^\Delta$ is equal to the \mathbb{F} -dimension of

$$\bigoplus_{(P,e_P) \in [\mathcal{F}_b]} \bigoplus_{\pi \in [N_G(P,e_P) \backslash \mathcal{P}_{(P,e_P)}(L,u) / \text{Aut}(L,u)]} \mathbb{F}\text{Proj}(ke_P C_G(P), u) \otimes_{\text{Aut}(L,u)_{\overline{(P,e_P,\pi)}}} V,$$

where $[\mathcal{F}_b]$ denotes a set of isomorphism classes of objects in \mathcal{F}_b , $\mathcal{P}_{(P,e_P)}$ is the set of group isomorphisms $\pi : L \rightarrow P$ with $\pi i_u \pi^{-1} \in \text{Aut}_{\mathcal{F}_b}(P, e_P)$, and $\text{Aut}(L, u)_{\overline{(P,e_P,\pi)}}$ is the stabilizer

in $\text{Aut}(L, u)$ of the G -orbit of (P, e_P, π) . Since b is a block with Frobenius inertial quotient, the block $kC_G(P)e_P$ is nilpotent for every nontrivial subgroup P of D , see for instance [L18, Theorem 10.5.2]. Therefore, we have $l(ke_PC_G(P)) = 1$, and hence the multiplicity formula reduces to

$$\bigoplus_{(P, e_P) \in [\mathcal{F}_b]} \bigoplus_{\pi \in [N_G(P, e_P) \backslash \mathcal{P}_{(P, e_P)}(L, u) / \text{Aut}(L, u)]} V^{\text{Out}(L, u)_{\overline{(P, e_P, \pi)}}}.$$

Let $\mathcal{Q}_{D \rtimes E, p}$ denote the set of pairs (P, s) of p -subgroups P of $D \rtimes E$ and p' -elements s of $N_{D \rtimes E}(P)$. Let also $[\mathcal{Q}_{D \rtimes E, p}]$ denote a set of representatives of $D \rtimes E$ -orbits on $\mathcal{Q}_{D \rtimes E, p}$ under the conjugation map. Recall from [BY22, Corollary 7.4] that the multiplicity of $S_{L, u, V}$ in $\mathbb{F}T_{D \rtimes E}^\Delta$ is equal to the \mathbb{F} -dimension of

$$\bigoplus_{\substack{(P, s) \in [\mathcal{Q}_{D \rtimes E, p}] \\ (\tilde{P}, \tilde{s}) \cong (L, u)}} V^{N_{D \rtimes E}(P, s)},$$

where for a pair $(P, s) \in \mathcal{Q}_{D \rtimes E, p}$ with $(\tilde{P}, \tilde{s}) \cong (L, u)$, we fix an isomorphism $\phi_{P, s} : L \rightarrow P$ with $\phi_{P, s}(^u l) = {}^s \phi_{P, s}(l)$ for all $l \in L$ and we view V as an $\mathbb{F}N_{D \rtimes E}(P, s)$ -module via the group homomorphism

$$N_G(P, s) \rightarrow \text{Out}(L, u) \quad (1)$$

that sends $g \in N_G(P, s)$ to the image of $\phi_{P, s}^{-1} \circ i_g \circ \phi_{P, s}$ in $\text{Out}(L, u)$.

(c) Let $\mathcal{P}_b(G, L, u)$ denote the set of triples (P, e, π) where $(P, e) \in \mathcal{F}_b$ and $\pi \in \mathcal{P}_{(P, e_P)}(L, u)$. Let also $\mathcal{Q}_{D \rtimes E, p}(L, u)$ denote the set of pairs (P, s) in $\mathcal{Q}_{D \rtimes E, p}$ with the property that $(\tilde{P}, \tilde{s}) \cong (L, u)$.

If $(P, e, \pi) \in \mathcal{P}_b(G, L, u)$, then $\pi i_u \pi^{-1} \in \text{Aut}_{\mathcal{F}_b}(P, e_P)$ by definition and since \mathcal{F}_b is equal to $\mathcal{F}_{D \rtimes E}(D)$, it follows that there exists a p' -element s of $N_{D \rtimes E}(P)$ with $\pi i_u \pi^{-1} = i_s$. This implies by [BY22, Lemma 3.3] that $(\tilde{P}, \tilde{s}) \cong (L, u)$ and therefore we have a map

$$\Psi : \mathcal{P}_b(G, L, u) \rightarrow \mathcal{Q}_{D \rtimes E, p}(L, u), \quad (P, e, \pi) \mapsto (P, s).$$

5.1 Lemma *The map Ψ induces a bijection*

$$\overline{\Psi} : [G \backslash \mathcal{P}_b(G, L, u) / \text{Aut}(L, u)] \rightarrow [\mathcal{Q}_{D \rtimes E, p}(L, u)].$$

Proof First we show that the map $\overline{\Psi}$ is well-defined. Let (P, e, π) and (Q, f, ρ) be two elements in $\mathcal{P}_b(G, L, u)$ that lie in the same $G \times \text{Aut}(L, u)$ -orbit. We need to show that $\overline{\Psi}(P, e, \pi) = \overline{\Psi}(Q, f, \rho)$. Write $\Psi(P, e, \pi) = (P, s)$ and $\Psi(Q, f, \rho) = (Q, t)$. Let $g \in G$ and $\varphi \in \text{Aut}(L, u)$ such that

$$g \cdot (P, e, \pi) \cdot \varphi = (Q, f, \rho).$$

Then (P, e) and (Q, f) lie in the same isomorphism class in $[\mathcal{F}_b]$ and hence P and Q are $D \rtimes E$ -conjugate since $\mathcal{F}_b = \mathcal{F}_{D \rtimes E}(D)$. Thus, there exists $h \in D \rtimes E$ with $i_g = i_h : P \rightarrow Q$. Hence $\rho = i_g \pi \varphi = i_h \pi \varphi : L \rightarrow Q$. Since $\varphi \in \text{Aut}(L, u)$, one has $\varphi \circ i_u = i_u \circ \varphi$. Therefore,

$$i_t = \rho i_u \rho^{-1} = i_h \pi \varphi i_u \varphi^{-1} \pi^{-1} i_{h^{-1}} = i_h \pi i_u \pi^{-1} i_{h^{-1}} = i_h i_s i_{h^{-1}} = i_{h s h^{-1}}.$$

This shows that $(Q, t) = h \cdot (P, s)$ and hence the map $\bar{\Psi}$ is well-defined.

Now we show that $\bar{\Psi}$ is surjective. Let $(P, s) \in \mathcal{Q}_{D \rtimes E, p}(L, u)$. Since $(\tilde{P}, \tilde{s}) \cong (L, u)$, again by [BY22, Lemma 3.3], there exists $\pi : L \rightarrow P$ such that $\pi i_u = i_s \pi$, i.e. $\pi i_u \pi^{-1} = i_s : P \rightarrow P$. Since $\mathcal{F}_{D \rtimes E}(D) = \mathcal{F}_b$, it follows that there exists $g \in N_G(P, e)$ with $i_s = i_g$, and hence $(P, e, \pi) \in \mathcal{P}_b(G, L, u)$ with $\bar{\Psi}(P, e, \pi) = (P, s)$. Thus, $\bar{\Psi}$ is surjective.

Finally, we show that $\bar{\Psi}$ is injective. Let $(P, e, \pi), (Q, f, \rho) \in \mathcal{P}_b(G, L, u)$ be elements with $\bar{\Psi}(P, e, \pi) = \bar{\Psi}(Q, f, \rho)$. Write $(P, s) = \Psi(P, e, \pi)$ and $(Q, t) = \Psi(Q, f, \rho)$. Then there exists $h \in D \rtimes E$ such that

$$h \cdot (P, s) = (Q, t).$$

Again, there exists $g \in G$ such that $i_g = i_h : P \rightarrow Q$. Define

$$\varphi := \pi^{-1} \circ i_g^{-1} \circ \rho : L \rightarrow L.$$

One has

$$\begin{aligned} \varphi \circ i_u &= \pi^{-1} \circ i_g^{-1} \circ \rho \circ i_u = \pi^{-1} \circ i_g^{-1} \circ i_t \circ \rho = \pi^{-1} \circ i_g^{-1} \circ i_g \circ i_s \circ i_g^{-1} \circ \rho \\ &= \pi^{-1} \circ i_s \circ i_g^{-1} \circ \rho = i_u \circ \pi^{-1} \circ i_g^{-1} \circ \rho = i_u \circ \varphi \end{aligned}$$

which shows that $\varphi \in \text{Aut}(L, u)$. Moreover, one has

$$g \cdot (P, e, \pi) \cdot \varphi = (Q, f, \rho)$$

and so the map $\bar{\Psi}$ is injective. □

5.2 Lemma *Let $(P, e, \pi) \in [G \backslash \mathcal{P}_b(G, L, u) / \text{Aut}(L, u)]$ and $(P, s) = \bar{\Psi}(P, e, \pi) \in [\mathcal{Q}_{D \rtimes E, p}]$. Then the image of $N_{D \rtimes E}(P, s)$ in $\text{Out}(L, u)$ is equal to $\text{Out}(L, u)_{\overline{(P, e, \pi)}}$.*

Proof We have $\pi i_u \pi^{-1} = i_s$ and hence the image of $N_{D \rtimes E}(P, s)$ is given by

$$\begin{aligned} N_{D \rtimes E}(P, s) &\rightarrow \text{Out}(L, u) \\ h &\mapsto \pi^{-1} \circ i_h \circ \pi \end{aligned}$$

Note that since ${}^h s = s$, we have $i_h i_s = i_s i_h$, i.e., $i_h \pi i_u \pi^{-1} = \pi i_u \pi^{-1} i_h$. Therefore the image is

$$\begin{aligned} \{\overline{\pi^{-1} i_h \pi} \mid h \in D \rtimes E, i_h : P \rightarrow P, {}^h s = s\} &= \{\overline{\pi^{-1} i_g \pi} \mid g \in N_G(P, e), i_g \pi i_u \pi^{-1} = \pi i_u \pi^{-1} i_g\} \\ &= \{\overline{\pi^{-1} i_g \pi} \in \text{Out}(L, u) \mid \pi^{-1} i_g \pi = i_g, g \in N_G(P, e)\} \\ &= \text{Out}(L, u)_{\overline{(P, e, \pi)}} \end{aligned}$$

as was to be shown. \square

Proof of Theorem 1.3: We need to show that for any $L \neq 1$, the multiplicities of a simple diagonal p -permutation functor $S_{L,u,V}$ in $\mathbb{F}T_{G,b}^\Delta$ and in $\mathbb{F}T_{D \rtimes E}^\Delta$ are equal. But this follows from Lemma 5.1 and Lemma 5.2. Now Part (i) follows from Theorem 1.2(i), and Part (ii) follows from [L18, Theorem 10.5.10]. \square

References

- [Br90] M. BROUÉ: Isométries parfaites, types de blocs, catégories dérivées. *Astérisque* No. **181-182** (1990), 61–92.
- [BP20] R. BOLTJE, P. PEREPELITSKY: p -permutation equivalences between blocks of group algebras. arXiv:2007.09253.
- [BX08] R. BOLTJE, B. XU: On p -permutation equivalences: between Rickard equivalences and isotypies *Trans. Amer. Math. Soc.* **360**(10) (2008) 5067–5087.
- [BY20] S. BOUC, D. YILMAZ: Diagonal p -permutation functors. *J. Algebra* **556** (2020), 1036–1056.
- [BY22] S. BOUC, D. YILMAZ: Diagonal p -permutation functors, semisimplicity, and functorial equivalence of blocks. *Adv. Math.* **411** (2022), 108799.
- [L18] M. LINCKELMANN: The block theory of finite group algebras. Vol. II. Cambridge University Press, Cambridge, 2018.
- [R96] J. RICKARD: Splendid equivalences: derived categories and permutation modules. *Proc. Lond. Math. Soc.* **72** (1996), 331–358.

Serge Bouc, CNRS-LAMFA, Université de Picardie, 33 rue St Leu, 80039, Amiens, France.
`serge.bouc@u-picardie.fr`

Deniz Yilmaz, Department of Mathematics, Bilkent University, 06800 Ankara, Turkey.
`d.yilmaz@bilkent.edu.tr`