

Méthodes de relaxation pour les équations d'Euler avec gravité

Vivien Desveaux



2 février 2015, Amiens
Séminaire A³ d'analyse

Joint work with:

- Markus Zenk (Würzburg)
- Christian Klingenberg (Würzburg)
- Christophe Berthon (Nantes)

Outline

- 1 Introduction
- 2 Relaxation models
- 3 Relaxation scheme and main properties
- 4 Numerical results

System of Euler equation with gravity

$$\begin{cases} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + p) = -\rho \partial_x \Phi \\ \partial_t E + \partial_x (E + p)u = -\rho u \partial_x \Phi \end{cases}$$

- ρ : density
- u : velocity
- $E = \rho e + \rho u^2/2$: total energy, with e the internal energy
- $\tau := 1/\rho$: specific volume
- $p = p(\tau, e)$: pressure given by a general law
- $\Phi(x)$: given smooth gravitational potential
example: constant gravitational field $\Phi(x) = gx$

System of Euler equation with gravity

$$\begin{cases} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + p) = -\rho \partial_x \Phi \\ \partial_t E + \partial_x (E + p)u = -\rho u \partial_x \Phi \end{cases}$$

- The system rewrites

$$\partial_t w + \partial_x f(w) = s(w) \partial_x \Phi,$$

with

- ▶ $w = (\rho, \rho u, E)^T$: vector of conservative variables
- ▶ $f(w) = (\rho u, \rho u^2 + p, (E + p)u)^T$: flux function
- ▶ $s(w) = (0, -\rho, -\rho u)^T$: source term
- Set of physical admissible states:

$$\Omega = \left\{ w \in \mathbb{R}^3, \quad \rho > 0, \quad e > 0 \right\}$$

Steady states at rest

General steady state:
$$\begin{cases} u = 0 \\ \partial_x p = -\rho \partial_x \Phi \end{cases}$$

- **Polytropic equilibrium:** $\Gamma \in (0, 1) \cup (1, +\infty)$

$$p = K \rho^\Gamma \quad \Leftrightarrow \quad \begin{cases} \rho = \left(\frac{\Gamma-1}{K\Gamma} (C - \Phi) \right)^{\frac{1}{\Gamma-1}} \\ p = K^{\frac{1}{1-\Gamma}} \left(\frac{\Gamma-1}{\Gamma} (C - \Phi) \right)^{\frac{\Gamma}{\Gamma-1}} \end{cases}$$

- **Isothermal equilibrium:** $\Gamma \rightarrow 1$

$$\begin{cases} \rho = e^{\frac{C-\Phi}{K}} \\ p = K e^{\frac{C-\Phi}{K}} \end{cases}$$

- **Incompressible equilibrium:** $\Gamma \rightarrow +\infty$

$$\begin{cases} \rho = \text{constant} \\ p + \rho \Phi = \text{constant} \end{cases}$$

Entropy

The pressure is assumed to satisfy the second law of thermodynamics
 $\Rightarrow$ existence of a specific entropy $s(\tau, e)$ which satisfies

$$-Tds = de + pd\tau, \quad \text{with } T > 0 \text{ the temperature}$$

Lemma

The smooth solutions satisfy the additional conservation laws

$$\partial_t \rho \mathcal{F}(s) + \partial_x \rho \mathcal{F}(s) u = 0,$$

for all smooth function $\mathcal{F}$.

Moreover the application $w \mapsto \rho \mathcal{F}(s)$ is strictly convex iff

$$\mathcal{F}'(s) > 0 \quad \text{and} \quad \frac{1}{c_p} \mathcal{F}'(s) + \mathcal{F}''(s) > 0.$$

Objectives

Derive a numerical scheme which has the following properties:

- Preservation of the set Ω
- Accurate approximation of all the steady states at rest
- Exact capture of the polytropic, isothermal and incompressible equilibria
- Discrete entropy inequalities
- General gravitational potential and general pressure law

Means

- Finite volume method (approximate Riemann solver)
- Relaxation method

References

- [Chalons et al. '10] : constant gravity field ($\Phi(x) = gx$)
- [Käpelli & Mishra '13] : isentropic steady states

- 1 Introduction
- 2 Relaxation models**
- 3 Relaxation scheme and main properties
- 4 Numerical results

Relaxation framework

Initial system

$$\partial_t w + \partial_x f(w) = 0$$

$$w \in \Omega \subset \mathbb{R}^d$$

Relaxation framework

Initial system

$$\partial_t w + \partial_x f(w) = 0$$

$$w \in \Omega \subset \mathbb{R}^d$$

Relaxation system

$$\partial_t W + \partial_x F(w) = \frac{1}{\varepsilon} R(W)$$

$$W \in \mathcal{O} \subset \mathbb{R}^N, \quad N > d$$

Relaxation framework

 Q


Initial system

$$\partial_t w + \partial_x f(w) = 0$$

$$w \in \Omega \subset \mathbb{R}^d$$


Relaxation system

$$\partial_t W + \partial_x F(w) = \frac{1}{\varepsilon} R(W)$$

$$W \in \mathcal{O} \subset \mathbb{R}^N, \quad N > d$$

- Matrix $Q \in M_{d,N}(\mathbb{R})$ s.t. $Q\mathcal{O} = \Omega$ and $QR(W) = 0, \quad \forall W \in \mathcal{O}$


Relaxation framework



- Matrix $Q \in M_{d,N}(\mathbb{R})$ s.t. $Q\mathcal{O} = \Omega$ and $QR(W) = 0, \quad \forall W \in \mathcal{O}$
- For all $w \in \Omega$, there exists a unique equilibrium $\mathcal{E}(w)$ s.t.

$$Q\mathcal{E}(w) = w, \quad R(\mathcal{E}(w)) = 0, \quad QF(\mathcal{E}(w)) = f(w)$$

Relaxation framework



- Matrix $Q \in M_{d,N}(\mathbb{R})$ s.t. $Q\mathcal{O} = \Omega$ and $QR(W) = 0, \quad \forall W \in \mathcal{O}$
- For all $w \in \Omega$, there exists a unique equilibrium $\mathcal{E}(w)$ s.t.

$$Q\mathcal{E}(w) = w, \quad R(\mathcal{E}(w)) = 0, \quad QF(\mathcal{E}(w)) = f(w)$$

- Equilibrium manifold : $\mathcal{M} \subset \mathcal{O} := \{\mathcal{E}(w), w \in \Omega\}$
- The relaxation system should be simpler: we need the exact solution of the Riemann problem

Relaxation scheme

Initially: approximation w_i^n at time t^n on the cell K_i

- ➊ **Going to the relaxation system** on the equilibrium manifold:

$$W_i^n = \mathcal{E}(w_i^n)$$

- ➋ **Time evolution** by the Godunov scheme for the system

$$\partial_t W + \partial_x F(W) = 0 \text{ (i.e. } \varepsilon = +\infty)$$

$$\rightarrow W_i^{n+1,-}$$

- ➌ **Relaxation** : Resolution of $\partial_t W = \frac{1}{\varepsilon} R(W)$ in the limit $\varepsilon = 0$

$$\rightarrow W_i^{n+1}$$

- ➍ **Return to the initial system** by projection on the equilibrium manifold:

$$w_i^{n+1} = \mathcal{Q} W_i^{n+1}$$

Remark : by multiplying $\partial_t W = \frac{1}{\varepsilon} R(W)$ by $\mathcal{Q}$, we get $\partial_t \mathcal{Q} W = 0$, so the update is given by $w_i^{n+1} = \mathcal{Q} W_i^{n+1,-}$

The Suliciu model ([Suliciu '98], [Bouchut '04]...)

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + \pi) = -\rho \partial_x \Phi \\ \partial_t E + \partial_x (E + \pi) u = -\rho u \partial_x \Phi \\ \partial_t \rho \pi + \partial_x (\rho \pi + a^2) u = \frac{\rho}{\varepsilon} (p(\tau, e) - \pi) \\ \partial_t \Phi = 0 \end{array} \right. \quad \begin{array}{l} \text{Equilibrium} \\ \pi = p(\tau, e) \end{array}$$

Eigenvalues: $u - \frac{a}{\rho}, \quad 0, \quad u, \quad u + \frac{a}{\rho}$

Advantage:

- All the fields are linearly degenerate

Difficulties:

- The order of the eigenvalues is not fixed *a priori*
- The Riemann invariants for the eigenvalue 0 are strongly nonlinear


Relaxation model with moving gravity

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + \pi) = -\rho \partial_x Z \\ \partial_t E + \partial_x (E + \pi) u = -\rho u \partial_x Z \\ \partial_t \rho \pi + \partial_x (\rho \pi + a^2) u = \frac{\rho}{\varepsilon} (p(\tau, e) - \pi) \\ \partial_t \rho Z + \partial_x \rho Z u = \frac{\rho}{\varepsilon} (\Phi - Z) \end{array} \right. \quad \begin{array}{l} \text{Equilibrium} \\ \pi = p(\tau, e) \\ Z = \Phi \end{array}$$

- Eigenvalues: $u - \frac{a}{\rho}$, u , $u + \frac{a}{\rho}$
- Fixed order of the eigenvalues: $u - \frac{a}{\rho} < u < u + \frac{a}{\rho}$
- All the fields are linearly degenerate
- There is a **missing Riemann invariant** in order to determine a unique Riemann solution
 $\Rightarrow$ We need a closure relation

Relaxation model with moving gravity

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + \pi) = -\rho \partial_x Z \\ \partial_t E + \partial_x (E + \pi) u = -\rho u \partial_x Z \\ \partial_t \rho \pi + \partial_x (\rho \pi + a^2) u = \frac{\rho}{\varepsilon} (p(\tau, e) - \pi) \\ \partial_t \rho Z + \partial_x \rho Z u = \frac{\rho}{\varepsilon} (\Phi - Z) \end{array} \right.$$



To approximate the equation

$$\partial_x p = -\rho \partial_x \Phi \quad \underset{\substack{\text{equilibrium} \\ \pi = p(\tau, e) \\ Z = \Phi}}{\Longleftrightarrow} \quad \partial_x \pi = -\rho \partial_x Z,$$

we propose the following closure relation:

$$\frac{\pi_R^* - \pi_L^*}{\Delta x} = -\bar{\rho}(\rho_L, \rho_R) \frac{Z_R - Z_L}{\Delta x},$$

where $\bar{\rho}$ is a suitable ρ -average (defined later)

Approximate Riemann solver

Relaxation equilibrium state:

$$\mathcal{E}(w) = (\rho, \rho u, E, \rho p(\tau, e), \rho \Phi)^T$$

Theorem

With the closure equation, the Riemann problem admits a unique solution $W_{\mathcal{R}}\left(\frac{x}{t}, W_L, W_R\right)$.

Moreover,

$$w^{eq}\left(\frac{x}{t}, w_L, w_R\right) := W_{\mathcal{R}}^{(\rho, \rho u, E)}\left(\frac{x}{t}, \mathcal{E}(w_L), \mathcal{E}(w_R)\right)$$

defines an approximate Riemann solver (in the sense of Harten, Lax and van Leer) for the Euler equations with gravity.

Complete relaxation reformulation

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + \pi) = -\bar{\rho}(X^-, X^+) \partial_x Z \\ \partial_t E + \partial_x (E + \pi) u = -\bar{\rho}(X^-, X^+) u \partial_x Z \\ \partial_t \rho \pi + \partial_x (\rho \pi + a^2) u = \frac{\rho}{\varepsilon} (p(\tau, e) - \pi) \\ \partial_t \rho Z + \partial_x \rho Z u = \frac{\rho}{\varepsilon} (\Phi - Z) \\ \partial_t X^- + (u - \delta) \partial_x X^- = \frac{1}{\varepsilon} (\rho - X^-) \\ \partial_t X^+ + (u + \delta) \partial_x X^+ = \frac{1}{\varepsilon} (\rho - X^+) \end{array} \right.$$

Equilibrium

$$\pi = p(\tau, e) \quad Z = \Phi \quad X^\pm = \rho$$

- For $\delta > 0$ small enough, the system is hyperbolic with eigenvalues

$$u - \frac{a}{\rho} < u - \delta < u < u + \delta < u + \frac{a}{\rho}.$$

- The system has a complete set of Riemann invariants.
- Leads to the same approximate Riemann solver $w^{eq}(\frac{x}{t}, w_L, w_R)$.

Cargo-LeRoux formulation ($\Phi(x) = gx$)

We introduce a potential $q = \int^x g\rho dx$. Then we have

$$\partial_t q = \int^x g \partial_t \rho dx = - \int^x g \partial_x \rho u dx = -g\rho u = -u \partial_x q.$$

So q is governed by

$$\partial_t \rho q + \partial_x \rho q u = 0.$$

Equivalent reformulation of the Euler equations with gravity:

$$\begin{cases} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + p) + \partial_x q = 0 \\ \partial_t E + \partial_x (E + p)u + u \partial_x q = 0 \\ \partial_t \rho q + \partial_x \rho q u = 0. \end{cases}$$

Relaxation model with the Cargo-LeRoux formulation

- Case $\Phi(x) = gx$ [Chalons et al. '10]

$$\begin{cases} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + \pi) + \partial_x q = 0 \\ \partial_t E + \partial_x (E + \pi) u + u \partial_x q = 0 \\ \partial_t \rho q + \partial_x \rho q u = 0 \\ \partial_t \rho \pi + \partial_x (\rho \pi + a^2) u = \frac{\rho}{\varepsilon} (p(\tau, e) - \pi) \end{cases} \quad \begin{array}{l} \text{Equilibrium} \\ \\ \\ \pi = p(\tau, e) \end{array}$$

- Extension for a general gravitational field


If we define the potential by $q = \int^x \rho \partial_x \Phi dx$, it no longer satisfies a transport equation. We enforce the natural relaxation model

$$\begin{cases} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + \pi) + \partial_x q = 0 \\ \partial_t E + \partial_x (E + \pi) u + u \partial_x q = 0 \\ \partial_t \rho q + \partial_x \rho q u = \frac{\rho}{\varepsilon} (\int^x \rho \partial_x \Phi dx - q) \\ \partial_t \rho \pi + \partial_x (\rho \pi + a^2) u = \frac{\rho}{\varepsilon} (p(\tau, e) - \pi) \end{cases} \quad \begin{array}{l} \text{Equilibrium} \\ \\ \\ \pi = p(\tau, e) \\ q = \int^x \rho \partial_x \Phi dx \end{array}$$

- 1 Introduction
- 2 Relaxation models
- 3 Relaxation scheme and main properties**
- 4 Numerical results

The relaxation scheme

w_i^n : approximation of the solution on the cell $(x_{i-1/2}, x_{i+1/2})$ at time t^n



CFL restriction

$$\frac{\Delta t}{\Delta x} \max_{i \in \mathbb{Z}} \left| u_i^n \pm \frac{a}{\rho_i^n} \right| \leq \frac{1}{2}$$

The update at time $t^{n+1} = t^n + \Delta t$ is defined by

$$\begin{aligned} w_i^{n+1} = & \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_i} w^{eq} \left(\frac{x - x_{i-1/2}}{\Delta t}, w_{i-1}^n, w_i^n \right) dx \\ & + \frac{1}{\Delta x} \int_{x_i}^{x_{i+1/2}} w^{eq} \left(\frac{x - x_{i+1/2}}{\Delta t}, w_i^n, w_{i+1}^n \right) dx \end{aligned}$$

Theorem (Robustness)

Assume the parameter a satisfies the following inequalities:

$$u_L - \frac{a}{\rho_L} < u^* < u_R + \frac{a}{\rho_R},$$

$$e_L + \frac{\pi_L^{*2} - p_L^2}{2a^2} > 0, \quad e_R + \frac{\pi_R^{*2} - p_R^2}{2a^2} > 0$$

Then the relaxation scheme preserves the set of physical states Ω .

Theorem (Well-balancedness for general steady states)

The relaxation scheme preserves exactly the initial data satisfying

$$\begin{cases} u_i^0 = 0, \\ \frac{p_{i+1}^0 - p_i^0}{\Delta x} = -\bar{\rho}(\rho_i^0, \rho_{i+1}^0) \frac{\Phi_{i+1} - \Phi_i}{\Delta x}. \end{cases}$$

Then $w_i^{n+1} = w_i^n$, $\forall i \in \mathbb{Z}$, $\forall n \in \mathbb{N}$.

Theorem (Exact preservation of the incompressible equilibrium)

If the initial data w_i^0 satisfies

$$u_i^0 = 0, \quad \rho_i^0 = M, \quad p_i^0 + \rho_i^0 \Phi_i = C,$$

then $w_i^n = w_i^0, \quad \forall n \in \mathbb{N}$.

Theorem (Exact preservation of the isothermal equilibrium)

Assume the average function $\bar{p}$ is defined by

$$\bar{p}(\rho_L, \rho_R) = \begin{cases} \frac{\rho_R - \rho_L}{\ln(\rho_R) - \ln(\rho_L)} & \text{if } \rho_L \neq \rho_R, \\ \rho_L & \text{if } \rho_L = \rho_R. \end{cases}$$

If the initial data w_i^0 satisfies

$$u_i^0 = 0, \quad \rho_i^0 = \exp\left(\frac{C - \Phi_i}{K}\right), \quad p_i^0 = K \exp\left(\frac{C - \Phi_i}{K}\right),$$

then $w_i^n = w_i^0, \quad \forall n \in \mathbb{N}$.

Theorem (Exact preservation of the polytropic equilibrium)

Assume the average function $\bar{\rho}$ is defined by

$$\bar{\rho}(\rho_L, \rho_R) = \begin{cases} \frac{\Gamma - 1}{\Gamma} \frac{\rho_R^\Gamma - \rho_L^\Gamma}{\rho_R^{\Gamma-1} - \rho_L^{\Gamma-1}} & \text{if } \rho_L \neq \rho_R, \\ \rho_L & \text{if } \rho_L = \rho_R. \end{cases}$$

If the initial data w_i^0 satisfies

$$\begin{cases} u_i^0 = 0, \\ \rho_i^0 = \left(\frac{\Gamma-1}{K\Gamma} (C - \Phi_i) \right)^{\frac{1}{\Gamma-1}}, \\ p_i^0 = K^{\frac{1}{1-\Gamma}} \left(\frac{\Gamma-1}{\Gamma} (C - \Phi_i) \right)^{\frac{\Gamma}{\Gamma-1}}, \end{cases}$$

then $w_i^n = w_i^0, \quad \forall n \in \mathbb{N}.$

Theorem

Assume the parameter a satisfy the following Whitham conditions:

$$a^2 > p(\tau_{L,R}, e_{L,R}) \partial_e p(\tau_{L,R}, e_{L,R}) - \partial_\tau p(\tau_{L,R}, e_{L,R}),$$

$$a^2 > p(\tau_{L,R}^*, e_{L,R}^*) \partial_e p(\tau_{L,R}^*, e_{L,R}^*) - \partial_\tau p(\tau_{L,R}^*, e_{L,R}^*).$$

Then the relaxation scheme satisfies

$$\rho_i^{n+1} \mathcal{F}(s_i^{n+1}) \leq \rho_i^n \mathcal{F}(s_i^n) - \frac{\Delta t}{\Delta x} \left(\{\rho \mathcal{F}(s) u\}_{i+1/2}^n - \{\rho \mathcal{F}(s) u\}_{i-1/2}^n \right)$$

for all smooth function $\mathcal{F}$ such that $w \mapsto \rho \mathcal{F}(s)$ is strictly convex.

The entropy numerical flux is defined by

$$\{\rho \mathcal{F}(s) u\}_{i+1/2}^n = F_{i+1/2}^\rho \times \begin{cases} \mathcal{F}(s_i^n) & \text{if } F_{i+1/2}^\rho > 0, \\ \mathcal{F}(s_{i+1}^n) & \text{if } F_{i+1/2}^\rho < 0, \end{cases}$$

with $F_{i+1/2}^\rho$ the ρ -component of the numerical flux.

Sketch of the proof (1)

- 1 We introduce

$$I(\pi, \tau) = \pi + a^2 \tau \quad \text{and} \quad J(\pi, e) = e - \frac{\pi^2}{2a^2}.$$

I and J are strong Riemann invariants, i.e they satisfy

$$\partial_t \Psi(I, J) + u \partial_x \Psi(I, J) = 0,$$

for all smooth function $\Psi : \mathbb{R}^2 \rightarrow \mathbb{R}$.

- 2 There exists functions $\bar{\tau}(I, J)$ and $\bar{e}(I, J)$ which satisfy

$$\bar{\tau} \left(I_{|\pi=p(\tau,e)}, J_{|\pi=p(\tau,e)} \right) = \tau \quad \text{and} \quad \bar{e} \left(I_{|\pi=p(\tau,e)}, J_{|\pi=p(\tau,e)} \right) = e$$

- 3 We define $\bar{s}(W) = s(\bar{\tau}(I, J), \bar{e}(I, J))$. This function reaches its minimum on the equilibrium manifold:

$$\bar{s}(W) \geq \bar{s} \left(W_{|\pi=p(\tau,e)} \right) = s \left(W_{|\pi=p(\tau,e)} \right).$$

Sketch of the proof (2)

- ④ $\bar{s}$ only depends on I and $J \Rightarrow \partial_t \rho \mathcal{F}(\bar{s}) + \partial_x \rho \mathcal{F}(\bar{s}) u = 0$.

We integrate this equation on $[x_{i-1/2}, x_{i+1/2}) \times [t^n, t^{n+1})$:

$$\begin{aligned} \int_{x_{i-1/2}}^{x_{i+1/2}} (\rho \mathcal{F}(\bar{s})) (W_{\Delta x}(x, t^{n+1})) dx &= \int_{x_{i-1/2}}^{x_{i+1/2}} (\rho \mathcal{F}(\bar{s})) (W_{\Delta x}(x, t^n)) dx \\ &\quad - \int_{t^n}^{t^{n+1}} (\rho \mathcal{F}(\bar{s}) u) (W_{\Delta x}(x_{i+1/2}, t)) dt \\ &\quad + \int_{t^n}^{t^{n+1}} (\rho \mathcal{F}(\bar{s}) u) (W_{\Delta x}(x_{i-1/2}, t)) dt. \end{aligned}$$

We deduce

$$\begin{aligned} \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} (\rho \mathcal{F}(\bar{s})) (W_{\Delta x}(x, t^{n+1})) dx &= \rho_i^n \mathcal{F}(s_i^n) \\ &\quad - \frac{\Delta t}{\Delta x} \left(\{\rho \mathcal{F}(s) u\}_{i+1/2}^n - \{\rho \mathcal{F}(s) u\}_{i-1/2}^n \right). \end{aligned}$$

Sketch of the proof (3)

⑤ By definition of the scheme:

$$\rho_i^{n+1} \mathcal{F}(s_i^{n+1}) = (\rho \mathcal{F}(s)) \left(\frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} W_{\Delta x}(x, t^{n+1})|_{\pi=p(\tau,e)} dx \right).$$

Jensen inequality ($w \mapsto \rho \mathcal{F}(s)$ is convex):

$$\rho_i^{n+1} \mathcal{F}(s_i^{n+1}) \leq \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} (\rho \mathcal{F}(s)) (W_{\Delta x}(x, t^{n+1})|_{\pi=p(\tau,e)}) dx.$$

$\mathcal{F}$ is increasing + the entropy is minimal on the equilibrium manifold:

$$\rho_i^{n+1} \mathcal{F}(s_i^{n+1}) \leq \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} (\rho \mathcal{F}(\bar{s})) (W_{\Delta x}(x, t^{n+1})) dx.$$

- 1 Introduction
- 2 Relaxation models
- 3 Relaxation scheme and main properties
- 4 Numerical results**

Isothermal equilibrium

- Computational domain: $[0, 1]$ until time $T = 0.25$

- $\Phi(x) = x^2$

- Initial condition:
$$\begin{cases} \rho_0(x) = e^{-x^2} \\ u_0(x) = 0 \\ p_0(x) = e^{-x^2} \end{cases}$$

- L^1 error:

N	Density	Velocity
100	1.01E-16	1.41E-16
200	2.24E-16	1.42E-16
400	3.61E-16	1.38E-16
800	5.39E-16	1.71E-16
1600	9.28E-16	2.49E-16
3200	1.60E-15	2.50E-16

General steady state

- Computational domain: $[0, 1]$ with periodic boundary conditions until time $T = 1$
- $\Phi(x) = -\sin(2\pi x)$
- Initial condition:
$$\begin{cases} \rho_0(x) = 3 + 2\sin(2\pi x) \\ u_0(x) = 0 \\ p_0(x) = 3 + 3\sin(2\pi x) - \frac{1}{2}\cos(4\pi x) \end{cases}$$

We check easily $\partial_x p_0 + \rho_0 \partial_x \Phi = 0$

- L^1 error:

N	Density		Velocity	
100	4.46E-05	—	2.03E-05	—
200	7.11E-06	2.65	5.29E-06	1.94
400	1.23E-06	2.53	1.34E-06	1.98
800	2.35E-07	2.39	3.37E-07	1.99
1600	5.02E-08	2.23	8.44E-08	2.00
3200	1.15E-08	2.13	2.11E-08	2.00


Perturbation of an isothermal equilibrium

- Computational domain: $[0, 1]$ until time $T = 0.25$

- $\Phi(x) = x$



- Initial condition:
$$\begin{cases} \rho_0(x) = e^{-x} \\ u_0(x) = 0 \\ p_0(x) = e^{-x} + 0.01e^{-100(x-0.5)^2} \end{cases}$$

- Pressure perturbation: $\delta p = p - p_0$



Perturbation of a polytropic equilibrium

- Computational domain: $[0, 2]$ until time $T = 1.5$
- $\Phi(x) = x$
- Initial condition:
$$\begin{cases} \rho_0(x) = (1 - 0.4\Phi(x))^{1.5} \\ u_0(x) = 0 \\ p_0(x) = (1 - 0.4\Phi(x))^{2.5} \end{cases}$$
- Velocity perturbation: $u(x=0) = 10^{-6} \sin(4\pi t)$



Perspectives

- Extension to second-order
 - ▶ MUSCL technique written as a convex combination of first-order scheme
 - ▶ Difficulty to preserve the well-balanced property
 - ▶ Possible solution: use a unusual set of variables for the reconstruction
- Extension to unstructured 2D
 - ▶ Convex combination of 1D scheme by interface