

# A high-order entropy satisfying scheme for the Euler equations

Christophe Berthon, Vivien Desveaux



# Introduction

- Hyperbolic system of conservation laws

$$\begin{cases} \partial_t w + \partial_x f(w) = 0 \\ w(x, 0) = w_0(x) \end{cases}$$

$w : \mathbb{R}^+ \times \mathbb{R} \rightarrow \Omega$  : unknown state vector

$f : \Omega \rightarrow \mathbb{R}^d$  : continuous flux function

$w_0 \in L^1_{\text{loc}}(\mathbb{R}; \Omega)$  : initial condition

- $\Omega \subset \mathbb{R}^d$  convex set of physical states
- Objective: study the stability of high-order space-time schemes

# Euler equations

$$\begin{cases} \partial_t \rho + \partial_x \rho u = 0 \\ \partial_t \rho u + \partial_x (\rho u^2 + p) = 0 \\ \partial_t E + \partial_x (E + p)u = 0 \end{cases}$$

- $\rho$ : density
- $u$ : velocity
- $E$ : total energy
- $p$ : pressure given by the perfect gas law

$$p = (\gamma - 1) \left( E - \frac{\rho u^2}{2} \right), \quad \gamma \in (1, 3]$$

- Set of physical states:

$$\Omega = \left\{ w \in \mathbb{R}^3, \rho > 0, p > 0 \right\}$$

# Weak solutions and entropy solutions

- A function  $w \in L^1_{\text{loc}}(\mathbb{R} \times \mathbb{R}^+; \Omega)$  is a **weak solution** if  $\forall \phi \in C^1_c(\mathbb{R} \times \mathbb{R}^+; \mathbb{R}^d)$ , we have

$$\int_{\mathbb{R}} \int_{\mathbb{R}^+} (w \cdot \partial_t \phi + f(w) \cdot \partial_x \phi) dt dx + \int_{\mathbb{R}} w(x, 0) \cdot \phi(x, 0) dx = 0.$$

- A convex function  $S \in C^2(\Omega; \mathbb{R})$  is an **entropy** for the system if there exists an entropy flux  $g \in C^2(\Omega; \mathbb{R})$  such that  $\nabla f(w) \nabla S(w) = \nabla g(w)$ ,  $\forall w \in \Omega$ .
- A weak solution  $w$  is an **entropy solution** if for any entropy pair  $(S, g)$  of the system, and  $\forall \phi \in C^1_c(\mathbb{R} \times \mathbb{R}^+; \mathbb{R})$ ,  $\phi \geq 0$ ,  $w$  satisfies

$$\int_{\mathbb{R}} \int_{\mathbb{R}^+} (S(w) \partial_t \phi + g(w) \partial_x \phi) dt dx + \int_{\mathbb{R}} S(w(x, 0)) \phi(x, 0) dx \geq 0.$$

- $(S, g)$  being an entropy pair of the system, an **entropy numerical flux** is a continuous function  $G : (\Omega)^{2s} \rightarrow \mathbb{R}$  which is consistent with  $g$ , i.e.  $G(w, \dots, w) = g(w)$ .

- 1 Motivations
- 2 Euler equations: from one to all discrete entropy inequalities
- 3 The E-MOOD scheme

# Space and time discretizations

- Space discretization: cells  $[x_{i-1/2}, x_{i+1/2}]$  with constant size  $\Delta x = x_{i+1/2} - x_{i-1/2}$
- Time discretization:  $t^n = n\Delta t$
- Rectangular cells in the  $(x, t)$ -plane:

$$R_i^n = [x_{i-1/2}, x_{i+1/2}) \times [t^n, t^{n+1})$$

- The sequence  $(\Delta x, \Delta t)$  of discretization steps is devoted to converge to  $(0, 0)$ , the ratio  $\frac{\Delta t}{\Delta x}$  being kept constant.

# A general high-order space-time scheme

## Initial condition

$$w_i^0 = \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} w_0(x) dx$$

## Runge-Kutta time discretization

$$w_i^{n,(\ell)} = w_i^n - \frac{\Delta t}{\Delta x} \sum_{j=0}^{\ell-1} c_{\ell,j} \left( F_{i+1/2}^{n,(j)} - F_{i-1/2}^{n,(j)} \right), \quad \ell = 1, \dots, m$$

$$w_i^{n,(0)} = w_i^n, \quad w_i^{n+1} = w_i^{n,(m)}$$

**Assumptions:**  $c_{\ell,j} \geq 0$ ,  $\sum_{j=0}^{m-1} c_{m,j} = 1$

## Space discretization

$$F_{i+1/2}^{n,(j)} = F \left( w_{i-s+1}^{n,(j)}, \dots, w_{i+s}^{n,(j)} \right)$$

**Assumptions:**  $F$  continuous and consistent ( $F(w, \dots, w) = f(w)$ )

# A general high-order space-time scheme

## Initial condition

$$w_i^0 = \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} w_0(x) dx$$

## Runge-Kutta time discretization

$$w_i^{n,(\ell)} = w_i^n - \frac{\Delta t}{\Delta x} \sum_{j=0}^{\ell-1} c_{\ell,j} \left( F_{i+1/2}^{n,(j)} - F_{i-1/2}^{n,(j)} \right), \quad \ell = 1, \dots, m$$

$$w_i^{n,(0)} = w_i^n, \quad w_i^{n+1} = w_i^{n,(m)}$$

We introduce the piecewise constant functions

$$w^\Delta(x, t) = w_i^n, \quad \text{for } (x, t) \in R_i^n,$$

$$w^{\Delta,(\ell)}(x, t) = w_i^{n,(\ell)}, \quad \text{for } (x, t) \in R_i^n.$$



# Lax-Wendroff Theorem

## Theorem

(i) Assume the following hypotheses:

- There exists a compact  $K \subset \Omega$  such that  $w^{\Delta,(\ell)} \in K$ ,  $\forall \ell = 0, \dots, m$ ;
- $w^{\Delta}$  converges in  $L^1_{loc}(\mathbb{R} \times \mathbb{R}^+; \Omega)$  to a function  $w$ .

Then  $w$  is a weak solution.

(ii) Assume the additional hypothesis:

- For all entropy pair  $(S, g)$ , there exists an entropy numerical flux  $G$ , such that we have the discrete entropy inequality (DEI)

$$\frac{S(w_i^{n+1}) - S(w_i^n)}{\Delta t} + \sum_{j=0}^{m-1} c_{m,j} \frac{G_{i+1/2}^{n,(j)} - G_{i-1/2}^{n,j}}{\Delta x} \leq 0,$$

$$\text{with } G_{i+1/2}^{n,(j)} = G(w_{i-s+1}^{n,(j)}, \dots, w_{i+s}^{n,(j)}).$$

Then  $w$  is an entropic solution.

# High-order time schemes

## Reformulation of the Runge-Kutta discretization (Shu-Osher)

$$w_i^{n,\ell} = \sum_{j=0}^{\ell-1} \left( \alpha_{\ell,j} w_i^{n,(j)} - \beta_{\ell,j} \frac{\Delta t}{\Delta x} \left( F_{i+1/2}^{n,(j)} - F_{i-1/2}^{n,(j)} \right) \right),$$

$$\text{with } \alpha_{\ell,j} > 0, \sum_{j=0}^{\ell-1} \alpha_{\ell,j} = 1.$$

**Assumption:**  $\beta_{\ell,j} > 0$ .

### Theorem

Assume the first-order time scheme  $w_i^{n+1} = w_i^n - \frac{\Delta t}{\Delta x} \left( F_{i+1/2}^n - F_{i-1/2}^n \right)$  satisfies the DEI  $\frac{S(w_i^{n+1}) - S(w_i^n)}{\Delta t} + \frac{G_{i+1/2}^n - G_{i-1/2}^n}{\Delta x} \leq 0$ , then the Runge-Kutta scheme satisfies the DEI

$$\frac{S(w_i^{n+1}) - S(w_i^n)}{\Delta t} + \sum_{j=0}^{m-1} c_{m,j} \frac{G_{i+1/2}^{n,(j)} - G_{i-1/2}^{n,(j)}}{\Delta x} \leq 0$$

# High-order space schemes

- No DEI like

$$\frac{S(w_i^{n+1}) - S(w_i^n)}{\Delta t} + \sum_{j=0}^{m-1} c_{m,j} \frac{G_{i+1/2}^{n,(j)} - G_{i-1/2}^{n,(j)}}{\Delta x} \leq 0$$

was ever proven as soon as the scheme is at least second-order in space.

- **Example:** second-order MUSCL scheme

- ▶ We consider  $L : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  a slope limiter and we define the limited slope  $\mu_i^{n,(j)} = L\left(w_i^{n,(j)} - w_{i-1}^{n,(j)}, w_{i+1}^{n,(j)} - w_i^{n,(j)}\right)$ .
- ▶ The MUSCL flux is defined by

$$F_{i+1/2}^{n,(j)} = F\left(w_i^{n,(j)} + \frac{1}{2}\mu_i^{n,j}, w_{i+1}^{n,(j)} - \frac{1}{2}\mu_{i+1}^{n,(j)}\right),$$

where  $F$  is a first-order numerical flux.

## DEI satisfied by the MUSCL scheme

The known DEI satisfied by the MUSCL scheme all write

$$\frac{S(w_i^{n+1}) - S(w_i^n)}{\Delta t} + \sum_{j=0}^{m-1} c_{m,j} \frac{G_{i+1/2}^{n,(j)} - G_{i-1/2}^{n,(j)}}{\Delta x} \leq \sum_{j=0}^{m-1} \alpha_{m,j} \frac{P_i^{n,(j)} - S(w_i^{n,(j)})}{\Delta t}$$

where  $P_i^{n,(j)} = P(w_i^{n,(j)}, \mu_i^{n,(j)}, \Delta x, S)$ .

- Examples of operator  $P$ :

$$P_1(w, \mu, \Delta x, S) = \frac{S(w - \mu/2) + S(w + \mu/2)}{2} \quad (\text{Berthon})$$

$$P_2(w, \mu, \Delta x, S) = \frac{1}{\Delta x} \int_{-\Delta x/2}^{\Delta x/2} S\left(w + \frac{x}{\Delta x} \mu\right) dx \quad (\text{Bouchut})$$

- The operator  $P$  satisfies:  $\exists C > 0$  such that

$$0 \leq P(w, \mu, \Delta x, S) - S(w) \leq C \|\nabla^2 S(w)\| \|\mu\|^2$$

## Convergence of $D^\Delta$ : theoretical study

- We define the piecewise constant function

$$D^\Delta(x, t) = \sum_{j=0}^{m-1} \alpha_{m,j} \frac{P_i^{n,(j)} - S(w_i^{n,(j)})}{\Delta t}, \quad \text{for } (x, t) \in R_i^n$$

- Let  $\mu$  be the weak-star limit of the sequence  $D^\Delta$ . Let  $\beta$  be the entropy dissipation measure defined as the weak-star limit of the sequence

$$b^\Delta(x, t) = \sum_{j=0}^{m-1} \alpha_{m,j} \frac{\|w_i^{n,(j)} - w_{i-1}^{n,(j)}\|^2}{\Delta x}, \quad \text{for } (x, t) \in R_i^n.$$


- $\mu$  is absolutely continuous with respect to  $\beta$ .

### Conjecture (Hou-le Floch)

The entropy dissipation measure  $\beta$  is concentrated on the curves of discontinuity of  $w$ .

# Numerical study: test cases

1-rarefaction



shock-shock




Figure: Exact solution in density

Figure: Exact solution in density

- $L^1$  error for the convergence:  $E^\Delta = \sum_i |\rho_i^N - \rho_{ex}(x_i, T)|$
- Convergence of  $D^\Delta$ :  $I^\Delta = \int_{[0,1] \times [0,T]} D^\Delta(x, t) dx dt$

# 1-rarefaction: convergence of first-order time schemes




Figure: Convergence of second-order space / first-order time schemes

# 1-rarefaction: what superbee does (1)




Figure: Solution given by the superbee limiter with 1000 cells



# 1-rarefaction: what superbee does (2)




Figure: Solution given by the superbee limiter with 2000 cells

# 1-rarefaction: convergence of $I^\Delta$ for first-order time schemes




Figure: Convergence of  $I^\Delta$  for second-order space / first-order time schemes

# 1-rarefaction: convergence of second-order time schemes




Figure: Convergence of second-order space-time schemes

# 1-rarefaction: convergence of $I^\Delta$ for second-order time schemes




Figure: Convergence of  $I^\Delta$  for second-order space-time schemes

# Shock-Shock: convergence of first-order time schemes




Figure: Convergence of second-order space / first-order time schemes

# Shock-Shock: convergence of second-order time schemes




Figure: Convergence of second-order space-time schemes

# Shock-Shock: convergence of $I^\Delta$ for second-order time schemes




Figure: Convergence of  $I^\Delta$  for second-order space-time scheme

# Conclusion

- Numerical results confirm the Hou-le Floch conjecture: when the scheme converges, the weak-star limit  $\mu$  of  $D^\Delta$  seems to be concentrated on the curves of discontinuity of  $w$ .
- This does not imply that the limit is not entropic, but only that the usual DEI are not the suitable tool to prove a Lax-Wendroff theorem.
- We have to focus on the stronger DEI

$$\frac{S(w_i^{n+1}) - S(w_i^n)}{\Delta t} + \sum_{j=0}^{m-1} c_{m,j} \frac{G_{i+1/2}^{n,(j)} - G_{i-1/2}^{n,j}}{\Delta x} \leq 0,$$

- Most of the limiters, when combined with a first-order scheme, seem to be unstable to small perturbations, though with a very low explosion rate.



- 1 Motivations
- 2 Euler equations: from one to all discrete entropy inequalities
- 3 The E-MOOD scheme

# The family of entropies for the Euler equations

The Euler system possesses a family of entropy pairs  $(S, g)$  written

$$S = -\rho h(s), \quad g = -\rho u h(s),$$

where  $s = \ln \left( \frac{p}{\rho^\gamma} \right)$  is the specific entropy and  $h$  is a smooth function satisfying

$$h'(s) > 0, \quad h'(s) - \gamma h''(s) > 0.$$

## Lemma (reformulation)

The entropy pairs of the Euler system write

$$S(r) = \rho \psi(r), \quad g(r) = \rho u \psi(r),$$

where  $r = \frac{\rho^{1/\gamma}}{p}$  and  $\psi$  is a smooth decreasing convex function.

We consider the scheme  $w_i^{n+1} = w_i^n - \frac{\Delta t}{\Delta x} (F_{i+1/2} - F_{i-1/2})$ , where  $w_i^n = (\rho_i^n, \rho_i^n u_i^n, E_i^n)^T$  and  $F_{i+1/2} = (F_{i+1/2}^\rho, F_{i+1/2}^{\rho u}, F_{i+1/2}^E)^T$ .

## Theorem

*Assume the scheme is  $\Omega$ -preserving. Assume the DEI*

$$-\rho_i^{n+1} r_i^{n+1} \leq -\rho_i^n r_i^n - \frac{\Delta t}{\Delta x} \left( -F_{i+1/2}^\rho r_{i+1/2}^n + F_{i-1/2}^\rho r_{i-1/2}^n \right)$$

*with  $r_{i+1/2}^n = \begin{cases} r_{i+1}^n & \text{if } F_{i+1/2}^\rho < 0 \\ r_i^n & \text{if } F_{i+1/2}^\rho > 0 \end{cases}$ . Assume the additional CFL like condition (Larrouturou)*

$$\frac{\Delta t}{\Delta x} \left( \max(0, F_{i+1/2}^\rho) - \min(0, F_{i-1/2}^\rho) \right) \leq \rho_i^n.$$

*Then the scheme satisfies all the discrete entropy inequalities.*

Example : the HLLC/Suliciu relaxation scheme

# Proof of the Theorem (1)

The numerical flux can be written

$$F_{i+1/2}^\rho r_{i+1/2} = F_{i+1/2}^\rho \frac{r_i^n + r_{i+1}^n}{2} - \left| F_{i+1/2}^\rho \right| \frac{r_{i+1}^n - r_i^n}{2}.$$

The DEI then writes

$$r_i^{n+1} \geq \frac{a}{\rho_i^{n+1}} r_{i-1}^n + \frac{b}{\rho_i^{n+1}} r_i^n + \frac{c}{\rho_i^{n+1}} r_{i+1}^n,$$

where we have set

$$\begin{aligned} a &= \frac{\Delta t}{2\Delta x} \left( F_{i-1/2}^\rho + \left| F_{i-1/2}^\rho \right| \right), \\ b &= \rho_i^n - \frac{\Delta t}{2\Delta x} \left( F_{i+1/2}^\rho + \left| F_{i+1/2}^\rho \right| - F_{i-1/2}^\rho + \left| F_{i-1/2}^\rho \right| \right), \\ c &= \frac{\Delta t}{2\Delta x} \left( \left| F_{i+1/2}^\rho \right| - F_{i+1/2}^\rho \right). \end{aligned}$$

## Proof of the Theorem (2)

- ▶ We have  $a + b + c = \rho_i^n - \frac{\Delta t}{\Delta x} \left( F_{i+1/2}^\rho - F_{i-1/2}^\rho \right) = \rho_i^{n+1}$ .
    - ▶  $a \geq 0, c \geq 0$
    - ▶  $b \geq 0$  thanks to the CFL like condition
- $\Rightarrow r_i^{n+1}$  is greater than a convex combination of  $r_{i-1}^n, r_i^n$  and  $r_{i+1}^n$ .
- We consider an entropy pair which can writes  $(S, g) = (\rho\psi(r), \rho u\psi(r))$  with  $\psi$  a smooth decreasing convex function thanks to the Lemma.
  - $\psi$  is decreasing:

$$\psi(r_i^{n+1}) \leq \psi\left(\frac{a}{\rho_i^{n+1}}r_{i-1}^n + \frac{b}{\rho_i^{n+1}}r_i^n + \frac{c}{\rho_i^{n+1}}r_{i+1}^n\right)$$

- Jensen inequality ( $\psi$  is convex):

$$\psi(r_i^{n+1}) \leq \frac{a}{\rho_i^{n+1}}\psi(r_{i-1}^n) + \frac{b}{\rho_i^{n+1}}\psi(r_i^n) + \frac{c}{\rho_i^{n+1}}\psi(r_{i+1}^n)$$

## Proof of the Theorem (3)

- We replace  $a$ ,  $b$  and  $c$  by their value to obtain

$$\begin{aligned}\rho_i^{n+1}\psi(r_i^{n+1}) &\leq \rho_i^n\psi(r_i^n) - \frac{\Delta t}{2\Delta x} \left( F_{i+1/2}^\rho(\psi(r_i^n) + \psi(r_{i+1}^n)) \right. \\ &\quad \left. - |F_{i+1/2}^\rho|(\psi(r_{i+1}^n) - \psi(r_i^n)) - F_{i-1/2}^\rho(\psi(r_{i-1}^n) + \psi(r_i^n)) \right. \\ &\quad \left. + |F_{i-1/2}^\rho|(\psi(r_i^n) - \psi(r_{i-1}^n)) \right).\end{aligned}$$

- We define  $\psi_{i+1/2}^n = \begin{cases} \psi(r_{i+1}^n) & \text{if } F_{i+1/2}^\rho < 0 \\ \psi(r_i^n) & \text{if } F_{i+1/2}^\rho > 0 \end{cases}$ .
- We have shown the DEI (for the entropy pair  $(\rho\psi(r), \rho u\psi(r))$ )

$$\rho_i^{n+1}\psi(r_i^{n+1}) \leq \rho_i^n\psi(r_i^n) - \frac{\Delta t}{\Delta x} \left( F_{i+1/2}^\rho\psi_{i+1/2}^n - F_{i-1/2}^\rho\psi_{i-1/2}^n \right).$$

□

- 1 Motivations
- 2 Euler equations: from one to all discrete entropy inequalities
- 3 The E-MOOD scheme

## First-order scheme

We consider a first-order scheme

$$w_i^{n+1} = w_i^n - \frac{\Delta t}{\Delta x} \left( F(w_i^n, w_{i+1}^n) - F(w_{i-1}^n, w_i^n) \right).$$

For a time step restricted according to the CFL condition

$$\frac{\Delta t}{\Delta x} \max_{i \in \mathbb{Z}} |\lambda^\pm(w_i^n, w_{i+1}^n)| \leq \frac{1}{2},$$

the first-order scheme is assumed to satisfy:

- (i)  $w_i^n \in \Omega, \quad \forall i \in \mathbb{Z} \quad \Rightarrow \quad w_i^{n+1} \in \Omega, \quad \forall i \in \mathbb{Z}$
- (ii)  $\forall i \in \mathbb{Z}$ , the following DEI is satisfied:

$$\begin{aligned} -\rho_i^{n+1} r_i^{n+1} \leq & -\rho_i^n r_i^n - \frac{\Delta t}{\Delta x} \left( -F^\rho(w_i^n, w_{i+1}^n) r_{i+1/2}^n \right. \\ & \left. + F^\rho(w_{i-1}^n, w_i^n) r_{i-1/2}^n \right). \end{aligned}$$

Example: HLLC scheme



# High-order reconstruction

- A reconstruction function is a continuous function  $\mathcal{R} : \Omega^{2s+1} \rightarrow \Omega$  such that  $\mathcal{R}(w, \dots, w) = w$ , for all  $w \in \Omega$ .
- A high-order reconstruction function is usually a reconstruction function based on high degree polynomial reconstruction.
- Here, we consider two reconstruction functions  $\mathcal{R}_-$  and  $\mathcal{R}_+$ . The associated MUSCL scheme is then given by

$$w_i^{n+1} = w_i^n - \frac{\Delta t}{\Delta x} (F(\mathcal{W}_{i,+}, \mathcal{W}_{i+1,-}) - F(\mathcal{W}_{i-1,+}, \mathcal{W}_{i,-})),$$

with  $\mathcal{W}_{i,\pm} = \mathcal{R}_{\pm}(w_{i-s}^n, \dots, w_{i+1}^n)$

- Example: second-order MUSCL scheme:

$$\mathcal{R}_{\pm}(w_{i-1}^n, w_i^n, w_{i+1}^n) = w_i^n \pm \frac{1}{2}L(w_i^n - w_{i-1}^n, w_{i+1}^n - w_i^n),$$

where  $L$  is a slope limiter.

# The E-MOOD algorithm

- ➊ **Evaluation of the reconstructed states.** The reconstructed states are given by  $\mathcal{W}_{i,\pm} = \mathcal{R}_{\pm}(w_{i-s}^n, \dots, w_{i+s}^n)$
- ➋ **Computation of the candidate solution  $w_i^*$ .** We compute a candidate solution  $w_i^*$  using the MUSCL scheme

$$w_i^* = w_i^n - \frac{\Delta t}{\Delta x} (F(\mathcal{W}_{i,+}, \mathcal{W}_{i+1,-}) - F(\mathcal{W}_{i-1,+}, \mathcal{W}_{i,-})).$$

- ➌ **DEI test.** If  $w_i^*$  does not satisfy the DEI test

$$-\rho_i^* r_i^* \leq -\rho_i^n r_i^n - \frac{\Delta t}{\Delta x} \left( -F_{i+1/2}^{\rho} r_{i+1/2}^n + F_{i-1/2}^{\rho} r_{i-1/2}^n \right),$$

with  $F_{i+1/2}^{\rho} = F^{\rho}(\mathcal{W}_{i,+}, \mathcal{W}_{i+1,-})$ , then we set  $\mathcal{W}_{i,\pm} = w_i^n$

- ➍ **Stopping criterion.**
  - ▶ If the DEI test is satisfied on all the cells, the candidate solution is valid and we set  $w_i^{n+1} = w_i^*$
  - ▶ else the solution is recomputed from step 2

# Stability and robustness of the E-MOOD scheme

## Theorem

*Assume the time step  $\Delta t$  is chosen in order to satisfy the two following CFL like conditions:*

$$\frac{\Delta t}{\Delta x} \max_{i \in \mathbb{Z}} (|\lambda^{\pm}(w_{i,+}, w_{i+1,-})|, |\lambda^{\pm}(w_{i,-}, w_{i,+})|) \leq \frac{1}{4}$$

$$\frac{\Delta t}{\Delta x} \left( \max(0, F_{i+1/2}^{\rho}) - \min(0, F_{i-1/2}^{\rho}) \right) \leq \rho_i^n.$$

*Then the E-MOOD method provides an updated solution  $w_i^{n+1}$  after a finite number of iterations. It is physically admissible, and it satisfies all the entropy inequalities.*

# 1-rarefaction: first-order time schemes




Figure: Convergence of first-order time schemes: E-MOOD vs MUSCL

# 1-rarefaction: second-order time schemes




Figure: Convergence of second-order time schemes: E-MOOD vs MUSCL

# Shock-Shock: first-order time schemes




Figure: Convergence of first-order time schemes: E-MOOD vs MUSCL

# Shock-Shock: second-order time schemes




Figure: Convergence of second-order time schemes: E-MOOD vs MUSCL

## Smooth problem

- $\rho_0(x) = \begin{cases} 1 & \text{if } x < 0.2 \text{ or } x > 0.8 \\ 1 + \exp\left(\frac{(x-0.5)^2}{(x-0.2)(x-0.8)}\right) & \text{if } 0.2 \leq x \leq 0.8 \end{cases}$   
 $u_0(x) = 1, p_0(x) = 1$
- Periodic boundary conditions




Figure: Initial and final solution in density for the smooth problem



# Smooth problem: convergence




Figure: Convergence: E-MOOD vs MUSCL

Thank you for your attention!!